

RESEARCH ARTICLE

Catalases Induction in High Virulence Pinewood Nematode *Bursaphelenchus xylophilus* under Hydrogen Peroxide-Induced Stress

Cláudia S. L. Vicente^{1,2}, Yoriko Ikuyo¹, Ryoji Shinya^{1,3}, Manuel Mota², Koichi Hasegawa^{1*}

1 Department of Environmental Biology, College of Bioscience & Biotechnology, Chubu University, Kasugai, Aichi, Japan, **2** ICAAM—Instituto de Ciências Agrárias e Ambientais Mediterrânicas, Departamento de Biologia, Universidade de Évora, Évora, Portugal, **3** HHMI and Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, California, United States of America

* koichihasegawa@isc.chubu.ac.jp



OPEN ACCESS

Citation: Vicente CSL, Ikuyo Y, Shinya R, Mota M, Hasegawa K (2015) Catalases Induction in High Virulence Pinewood Nematode *Bursaphelenchus xylophilus* under Hydrogen Peroxide-Induced Stress. PLoS ONE 10(4): e0123839. doi:10.1371/journal.pone.0123839

Academic Editor: Jeum Kyu Hong, Gyeongnam National University of Science and Technology, KOREA, REPUBLIC OF

Received: September 3, 2014

Accepted: February 24, 2015

Published: April 20, 2015

Copyright: © 2015 Vicente et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by the Japan Society for Science Promotion (JSPS - reference P14394) postdoctoral fellowship to CSLV; the Heiwa Nakajima Foundation, international joint research grant; the European Project REPHRAME - Development of improved methods for detection, control and eradication of pine wood nematode in support of EU Plant Health policy, European Union Seventh Framework Programme FP7-KBBE-2010-4;

Abstract

Considered an EPPO A2 quarantine pest, *Bursaphelenchus xylophilus* is the causal agent of the pine wilt disease and the most devastating plant parasitic nematode attacking coniferous trees in the world. In the early stages of invasion, this nematode has to manage host defence mechanisms, such as strong oxidative stress. Only successful, virulent nematodes are able to tolerate the basal plant defences, and furthermore migrate and proliferate inside of the host tree. In this work, our main objective was to understand to what extent *B. xylophilus* catalases are involved in their tolerance to oxidative stress and virulence, using as oxidant agent the reactive oxygen species hydrogen peroxide (H₂O₂). After 24 hours of exposure, high virulence isolates of *B. xylophilus* could withstand higher H₂O₂ concentrations in comparison with low virulence *B. xylophilus* and *B. mucronatus*, corroborating our observation of *Bxy-ctl-1* and *Bxy-ctl-2* catalase up-regulation under the same experimental conditions. Both catalases are expressed throughout the nematode intestine. In addition, transgenic strains of *Caenorhabditis elegans* overexpressing *B. xylophilus* catalases were constructed and evaluated for survival under similar conditions as previously. Our results suggest that catalases of high virulence *B. xylophilus* were crucial for nematode survival under prolonged exposure to *in vitro* oxidative stress, highlighting their adaptive response, which could contribute to their success in host conditions.

Introduction

From 1905 until the present time, pine wilt disease (PWD) has been one of the most threatening diseases for worldwide forests [1]. PWD is considered a complex disease resulting from the interaction of three main elements: the pathogenic agent *Bursaphelenchus xylophilus* (PWN, pine wood nematode), the insect-vector *Monochamus* spp. (responsible for PWN dissemination),

and FEDER Funds through the Operational Programme for Competitiveness Factors - COMPETE and National Funds through FCT - Foundation for Science and Technology under the Strategic Project PEst-C/AGR/UI0115/2011. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

and the host tree, mostly from *Pinus spp.* [2–3]. In a *B. xylophilus* population, there is intra-specific variability in the ability to cause PWD (virulence) [4]. The existence of different levels of virulence among PWN isolates, and their relation to PWN reproductive ability *in vitro* and *in vivo* were already reported [5–6]. In this context, the low virulence nematodes, often designated as avirulent, show lower reproduction rates and increased generation times that eventually affect their host invasion and colonization [7–9].

In the early stage of PWD, PWN has to cope with different levels of plant immune responses. The first, non-specific, host tree reaction to nematode invasion (or other biotic/abiotic stress) is oxidative burst, an excessive production of reactive oxygen species (ROS). The predominant ROS in plant oxidative burst is H_2O_2 [10–11]. H_2O_2 is relatively stable, cell wall diffusible, and found to be transversal in different plant-pathogens systems, being a fundamental signal for inducing plant resistance (*i.e.* involved in cell-wall reinforcement or induction of defence-related genes in healthy adjacent tissues) [12]. H_2O_2 is particularly important since it is extremely reactive and can lead to the formation of radical hydroxyl OH^\cdot , for which no specific scavenger exists [12]. Generally, parasitic nematodes respond to an oxidant threat by increasing their protective antioxidant enzyme levels [13]. The major antioxidant enzyme families considered are: superoxide dismutases (SOD, EC 1.15.1.1) for conversion of superoxide anion to H_2O_2 , glutathione peroxidases (GPX, EC 1.11.1.9), catalases (CTL, E.C. 1.11.1.6) and peroxiredoxins (PRDX, EC 1.11.1.15) for the conversion of H_2O_2 into water [14]. Furthermore, many parasites employ many enzyme groups under oxidative stress conditions [14]. In plant parasitic nematodes, Molinari and Miacola [15] showed the presence of many antioxidant enzymes and the relation to its lifestyle stages of *Meloidogyne incognita* and *M. hapla*, *Globodera rostochiensis* and *G. pallida*, *Heterodera schachtii* and *H. carotae*, and *Xiphinema index*. Robertson et al. [16] described an important role of PRDX and GPX in *G. rostochiensis*. Bellafiore et al. [17] reported the presence of several detoxifying enzymes, in particular several glutathione S-transferases (GST, EC 2.5.1.18) in the secretome of *M. incognita* as means for controlling the global oxidative status and potential nematode virulence. PWN also has an efficient antioxidant system to reduce the deleterious effects of tree oxidative burst [18], mostly present in the nematode secretome [19] and expressed in the cuticle [20].

Pine oxidative burst was found intrinsically related with the production of ROS in the first stages of *B. xylophilus* invasion [21–22]. He et al. [23] evaluated the content of H_2O_2 of *P. thunbergii*, *P. massoniana* and *P. taeda* at the early stage of *B. xylophilus* invasion, and found that H_2O_2 content was higher in the susceptible pines than the resistant *P. taeda*. In this study, we specifically analyzed the tolerance of *B. xylophilus* isolates with different levels of virulence (high and low virulence) under *in vitro* oxidative stress conditions by assessing the effects of H_2O_2 exposure to nematode survival, catalase activity, catalase gene expression and their spatial localization. Additionally, we expressed *B. xylophilus* in the nematode model *C. elegans* to explore the efficiency of *B. xylophilus* catalases in promoting survival.

Material and Methods

B. xylophilus isolates and culturing

Two *B. xylophilus* isolates with high virulence (Bx Ka4 and Bx T4) and one isolate with low virulence (Bx C14-5) [6] were used in this study. In addition, one isolate of *B. mucronatus* (Un1), considered a low virulence nematode, was also used. All isolates were supplied by the FFPRI (Forestry and Forest Products Research Institute). The origin of nematode isolates Bx Ka4 and Bx C14-5 is described by Aikawa and Kikuchi [6]. The isolate Bx T4 was collected from dead

Pinus densiflora in 1992, in Iwate prefecture, Japan and *B. mucronatus* Un1 was obtained from *Monochamus alternatus*, in the Kyoto prefecture.

Nematodes were cultured in *Botrytis cinerea* on autoclaved barley seed at 25°C. Prior to experiments, nematodes were extracted overnight using Baermann funnel technique at 25°C. Nematodes were washed three times with sterilized distilled water with 10 min. centrifugations at 1,000 rpm between washes, surface cleaned with 3% L-lactic acid for 30s, and washed once with sterilized distilled water [9]. Mix-stages nematodes were used in all experiments.

Oxidative stress tolerance test

H₂O₂ was used as oxidative agent at concentrations ranging between 15 and 40 mM for *Bursaphelenchus* isolates. After nematode surface sterilization, the concentration of *Bursaphelenchus* was adjusted to 150 nematodes per 50 µl of sterile water. Mortality of nematodes was scored after 24h-stress exposure. Nematodes were considered dead if no movements were observed after mechanical stimulation. This experiment was repeated three independent times, with two technical replications for each.

H₂O₂ neutralization

H₂O₂ neutralization was inferred using an H₂O₂ Assay Kit (abcam, Massachusetts, USA). For this purpose, 150 nematodes were exposed for 24h in 15 mM H₂O₂. Afterwards, nematodes were pelleted at 1,000 g for 15 min and the supernatant was used for analysis following the manufacturer's protocol. This assay was repeated three independent times with two technical replications.

Relative gene expression of catalases under oxidative stress

Relative gene expressions of CTL (catalase) enzymes (*Bxy-ctl-1* and *Bxy-ctl-2*), previously predicted by Vicente et al. [24], were analyzed by qRT-PCR using SYBR green assay. Total RNA was extracted from 24h-stressed nematodes (approximately 2,500) in 15 mM H₂O₂ using CelLamp Direct RNA Prep Kit for RT-PCR (Takara Bio Inc., Japan) and following manufacturer's instructions. The concentration was quantified using NanoVue plus spectrophotometer (GE Healthcare Life Sciences, USA). Total RNA (adjusted to a concentration of 50 ng/µl) was reverse transcribed using oligo dT primer and PrimeScript RT enzyme from PrimeScript RT reagent Kit (Perfect Real Time, Takara Bio Inc., Japan). qRT-PCR was performed using CFX96 Real-Time (Bio-Rad Laboratories, Inc., California), and SYBR Premix Ex Taq II (Tli RnaseH Plus) kit (Takara Bio Inc., Japan). Actin (*Bxy-act-1*) was used as a reference gene for calculating relative expression levels of CTLs genes [25]. Primers were designed using Prime 3 software [26] and tested for specificity prior to qRT-PCR (Table 1). For each treatment, two experimental replications were conducted with two technical replications each. Controls with no template added were prepared for each qPCR run. Thermal cycling conditions were: initial denaturation at 95°C for 30 sec; 39 cycles of denaturation at 95°C for 5 sec, annealing and extension at 60°C for 30 sec; followed by the melting curve. A single peak at the melting temperature of the PCR-product confirmed primer specificity.

In-situ hybridization

Single-stranded DNA probes (sense and antisense) labelled with digoxigenin (Roche, United States of America) were used to detect mRNA in mix-stage *B. xylophilus* as described by De Boer et al. [28]. Table 1 presents the primers used to generate Bxy-CTL-1 and Bxy-CTL-2 probes (Table 1). As positive control for successful hybridization, ENG (cellulase) probe was

Table 1. Characterization of *Bursaphelenchus xylophilus* CTLs, and primers used for CTLs expression analysis.

Gene and Protein DB	ORF ¹	MW (Da)	SignalP ²	Secretome ³ (Sec/whole)
<i>Bxy-ctl-1</i> (BUX.s00579.159)	513	58836.42	NO	NO (ND)
<i>Bxy-ctl-2</i> (BUX.s01109.377)	272	54869.71	NO	YES (0.26)
Primers for qRT-PCR (5'– 3')				
<i>Bxy-ctl-1</i>	For: GCCAGCGTCTTCAGCAAAGT; Rev: CCAAATTCGTCATCGGTGT			
<i>Bxy-ctl-2</i>	For: CCGACTTCTTTCAACGGAAC; Rev: CCTTCATCGAGCACCTTTTC			
<i>Bxy-act-1</i>	For: CATCCTCCGTCTCGACTTGG; Rev: ATGTCACGCACGATTTACAG			
Primers for <i>In-situ</i> hybridization (5'– 3')				
<i>Bxy-CTL-1</i>	For: CTTCCGATCAACTGCCCTT; Rev: ATTTAGACAAGGGGCCAGCC			
<i>Bxy-CTL-2</i>	For: AGAGACCCCATCCTCTTCCC; Rev: AGCTTCTCCCTTTGCGTTGA			

Sec/whole—Normalized secreted protein/whole proteins; ND—not detected.

¹ ORF, open reading frame.

²Petersen et al. [27].

³ Shinya et al. [19].

doi:10.1371/journal.pone.0123839.t001

also prepared as described by Kikuchi et al. [29]. Images were acquired on an Olympus Bx50 light microscope at 10x-40x magnifications.

Transgenic *C. elegans* construction

C. elegans culturing and handling were carried out at 20°C as described by Brenner [30]. Strains used in this experiments were N2 (Bristol strain) and DP38 *unc-119(ed3)III*. To make reporter constructs, all PCRs were performed with Takara PrimeSTAR GXL DNA polymerase (Takara Bio Inc., Japan). *B. xylophilus* *ctl-1* and *-2* cDNAs were fused with *C. elegans* *ctl-1* or *-3* promoters and ligated into the *gfp* vector pPD95.77 (kindly provided by A. Fire, Stanford University) with In-Fusion HD Cloning system (Takara Bio Inc., Japan). We cloned a 1,940 bp sequence upstream from the *Ce-ctl-3* start codon for *Ce-Pctl-3*, and a 1,600 bp sequence between *Ce-ctl-3* and *Ce-ctl-1* CDS (coding sequences) for *Ce-Pctl-1*. Fusion genes created in this experiments were *Pctl-1::Bxy-ctl-1::gfp* and *Pctl-3::Bxy-ctl-2::gfp*, and PCR primer sequences are listed in Table 2. Each reporter construct (100 µg/ml) was co-injected with an equal concentration of pDP#MM016B into the gonadal arms of *unc-119(ed3)* adult hermaphrodites to obtain KHA149 {*unc-119(ed3)III*; *chuEx149[Pctl-1::Bxy-ctl-1::gfp, pDP#MM016B]*}, KHA150 {*unc-119(ed3)III*; *chuEx150[Pctl-1::Bxy-ctl-1::gfp, pDP#MM016B]*}, KHA151 {*unc-119(ed3)III*; *chuEx151[Pctl-3::Bxy-ctl-2::gfp, pDP#MM016B]*}, and KHA152 {*unc-119(ed3)III*; *chuEx152[Pctl-3::Bxy-ctl-2::gfp, pDP#MM016B]*}. Successful transgenic *C. elegans* were further confirmed by visualization of

Table 2. List of primers used in *C. elegans* constructs.

Genes	Primer 5'–3'
<i>Cectl-1Prom_lfs</i>	For: CGACTCTAGAGGATCATTGTTTGATATTCAAACCTTTTGTGTA; Rev: CGTCATTTTGGTTCTGAAATTTTAGTTAGG
<i>Cectl-3Prom_lfs</i>	For: CGACTCTAGAGGATCCTTCGTCACACTTCTATGGAATCC; Rev: GCCATTTTGAAGATTTACTGTTGAATTTCCG
<i>Bxy-ctl-1_lfs</i>	For: CAAACCAAAATGACGGACTTTGCGGCCAATC; Rev: CCAATCCCGGGGATCTTGACATCCTCTTGCAATCTCCAC
<i>Bxy-ctl-2_lfs</i>	For: AATCTTCAAATGGCAAACAAGAAGACTGC; Rev: CCAATCCCGGGGATCTCAAATGGGCCTTGACCTTCTTGG

doi:10.1371/journal.pone.0123839.t002

fluorescence expression patterns with a Nikon SMZ800 dissection microscope equipped with a fluorescence filter and a ZEISS LSM710 confocal laser-scanning microscope.

Oxidative stress tolerance in *C. elegans*

Synchronized L1 stage *C. elegans* were obtained by treating egg-containing/or gravid adults with sodium hypochlorite [30] and allowed to grow on NGM plates seeded with *E. coli* OP50 at 20°C for 48 hours until the late L4 stage. L4 stage *C. elegans* were washed with M9 buffer and adjusted to 20 nematodes per 50 µl of M9 buffer. Nematodes were transferred to a 96-well plate containing each concentration of H₂O₂ (0 to 250 µM). The plate was incubated at 20°C for 24 hours, and then examined for animal viability under a stereomicroscope. Since our transgenic nematodes used in this experiment retained transgenes as an extrachromosomal DNA array, non-transgenic uncoordinated nematodes appeared at a certain proportion (around 30%). We excluded non-transgenic uncoordinated animals when performing this experiment. All tests were carried out at 20°C. This experiment was repeated three independent times, and in each experimental repetition, two technical replications were used.

Data analysis

Statistical analysis was performed using SPSS version 11.5. One-way ANOVA analysis was performed in order to test if the different *B. xylophilus* and *B. mucronatus* isolates were equally tolerant to H₂O₂ treatments for the parameters: nematode survival percentage, H₂O₂ neutralization, and *C. elegans* survival percentage. *Post hoc* multi-comparison Duncan test (significance level of 0.05) was used to group significantly different *Bursaphelenchus* sp. isolates in the following parameters: nematode survival percentage, H₂O₂ neutralization, and *C. elegans* survival percentage. Levene's Test of Homogeneity was performed a priori to infer ANOVA assumption of homogeneity of variance.

Relative gene expressions of CTLs were analysed using the 2^{-ΔΔCT} method [25]. The data were analysed with C_T (cycle threshold) values in normal and stress conditions and using Eq. (1), where $\Delta\Delta C_T = (C_{T,Target} - C_{T,Actin})_{normal} - (C_{T,Target} - C_{T,Actin})_{stress}$. The fold change of CTL genes, normalized to *β-actin* and relative to the expression at normal conditions, were calculated for each sample using Eq. (1). Data represent the mean ± standard deviation (SD). Statistical differences at 95% and 99% confidence levels were inferred by Student t-test by comparison between relative expression of both CTLs for each *B. xylophilus* isolate and the control treatment without H₂O₂.

Results

Oxidative stress tolerance and H₂O₂ neutralisation

B. xylophilus (high virulence: Bx Ka4 and Bx T4; low virulence: Bx C14-5) and *B. mucronatus* (low virulence: Bm Un1) isolates were tested in increasing concentrations of H₂O₂ (0, 15, 20, 30 and 40 mM). After a 24h-exposure to this oxidant, nematode survival was examined (Fig 1). A clear difference between high and low virulence isolates was observed, even in the lowest H₂O₂ concentration (15 mM). The isolate Bx Ka4 seemed to be the most tolerant to H₂O₂ at all concentrations followed by Bx T4. Statistical differences ($p < 0.05$) between Bx Ka4 and Bx T4 were found in all H₂O₂ concentrations. Low virulence isolates (Bm Un1 and Bx C14-5) were sensitive to H₂O₂, having low survival even in 15 mM H₂O₂. No statistical differences ($p > 0.05$) were found between isolates.

In terms of H₂O₂ neutralization (Fig 2), high virulence Bx Ka4 and Bx T4 were more efficient at H₂O₂ reduction ($p < 0.05$) in comparison with the low virulence Bm Un1 and Bx C14-5 (no statistical differences between each pair).




Fig 1. Survival percentage of *Bursaphelenchus* sp. isolates (high virulence *B. xylophilus* Ka4 and *B. xylophilus* T4, and low virulence *B. xylophilus* C14-5 and *B. mucronatus* Un1) after 24 hours exposition to H₂O₂ conditions (H₂O₂ concentrations ranging from 0 to 40 mM). Error bars represent standard deviation. Different letters above the columns indicate significant differences ($p < 0.05$) between *Bursaphelenchus* sp. isolates survival percentages in each H₂O₂ treatment, according to *post-hoc* Duncan's test.

doi:10.1371/journal.pone.0123839.g001




Fig 2. H₂O₂ neutralization of *Bursaphelenchus* sp. isolates (high virulence *B. xylophilus* Ka4 and *B. xylophilus* T4, and low virulence *B. xylophilus* C14-5 and *B. mucronatus* Un1) after 24 hours exposition to H₂O₂. Error bars represent standard deviation. Different letters above the columns indicate significant differences ($p < 0.05$) between *Bursaphelenchus* sp. isolates H₂O₂ neutralization, according to *post-hoc* Duncan's test.

doi:10.1371/journal.pone.0123839.g002




Fig 3. Relative gene expression of *Bxy-ctl-1* (A) and *Bxy-ctl-2* (B) of *B. xylophilus* (high virulence isolates Ka4 and T4; low virulence isolate C14-5) after 24 hours exposition to H_2O_2 . * and ** indicate, respectively, statistical differences at 95% and 99% confidence levels compared to a normalized value of 1.00 for control treatment without H_2O_2 . Error bars represent standard deviation.

doi:10.1371/journal.pone.0123839.g003

Gene expression analysis

B. xylophilus catalases (*Bxy-ctl-1* and *Bxy-ctl-2*; Table 1) were previously analyzed and found to be conserved among Nematoda [24]. The CDS of both CTLs were similar between high and low virulence *B. xylophilus* isolates (data not shown). Because the lack of *B. mucronatus* genome in databases prevented the analyses of gene expression for the Bm Un1 isolate, our studies about levels of virulence will only focus on the Bx Ka4, Bx T4 and Bx C14-5 isolates. Relative gene expression of the CTLs was analyzed after a 24h-exposure to 15 mM H_2O_2 (Fig 3). High virulence isolates Bx Ka4 and Bx T4 presented a higher induction of CTL expression than low virulence Bx C14-5. In fact, the relative expression of both *Bxy-ctl-1* and *Bxy-ctl-2* of Bx C14-5 in stress conditions was not significantly different ($p > 0.05$) from expression in non-stress conditions. In oxidative conditions, *Bxy-ctl-1* and *Bxy-ctl-2* from both high virulence isolates Bx Ka4 and Bx T4 were significantly induced ($0.01 < p < 0.05$) in comparison with non-stress conditions (Fig 3A and 3B). Both *Bxy-ctl-1* and *Bxy-ctl-2* of Bx Ka4 were 2.5-fold up-regulated. *Bxy-ctl-1* of Bx T4 was significantly induced, nearly 2-fold (Fig 3A). *Bxy-ctl-2* was also significantly up regulated ($p < 0.05$) in stress, nearly 4-fold more than in normal conditions (Fig 3B).

Bxy-ctl-1 and *Bxy-ctl-2* messenger RNAs presented the same expression pattern in the nematode's intestine (Fig 4A and 4B). *Bxy-eng-1*, profusely expressed in *B. xylophilus* oesophageal glands, was used as a positive control (Fig 4D) [29]. No hybridization was observed for the control sense probe of *Bxy-ctl-1* (Fig 4C), *Bxy-ctl-2* or *Bxy-egn-1* (data not shown).

Transgenic *C. elegans* and OS tolerance

Three catalase genes exist in tandem in the *C. elegans* genome (*Ce-ctl-3*, -1, and -2), while two genes exist separately in the *B. xylophilus* genome (WormBase, <http://www.wormbase.org/>). Because *Bxy-ctl-1* and *Bxy-ctl-2* sequences were highly homologous to *Ce-ctl-1* and *Ce-ctl-3* respectively, and *Ce-Pctl-1* is identical to *Ce-Pctl-2*, we constructed transgenic *C. elegans* overexpressing *Bx-ctl-1* or *Bxy-ctl-2* with *C. elegans* promoters *Ce-Pctl-1* or *Ce-Pctl-3*. We created two independent transgenic lines for each construct; KHA149 and 150 for *Ce-Pctl-1::Bx-ctl-1::gfp*, and KHA151 and 152 for *Ce-Pctl-3::Bx-ctl-2::gfp*.

Under the control of *C. elegans* promoters, *B. xylophilus* *Bxy-ctl-1* and *Bxt-ctl-2* showed different spatial expression patterns; *ctl-1* was mainly detected in the cytosol of the intestine, and *ctl-2* was detected as high levels along the nervous system and pharynx (Fig 5A and 5B). These expression patterns were the same with those previously reported [31]. Although these




Fig 4. mRNA expression patterns of *Bxy-ctl-1* (A), *Bxy-ctl-2* (B), and *Bxy-eng-1* (D) in *B. xylophilus* Ka4. No expression signal was observed with *Bxy-ctl-1* sense probe (C). Light microscope images of *B. xylophilus* head region. Scale bars, 100 μ m.

doi:10.1371/journal.pone.0123839.g004


transgenic *C. elegans* were extrachromosomal lines with unstable transgene transmission, the expression patterns were similar between the transgenic lines KHA149 and 150, and between KHA151 and 152 (data not shown). We used KHA149 and 151 for further experiments.

To gauge *C. elegans* (wild-type and transgenic) tolerance to oxidative stress conditions, they were exposed to H_2O_2 concentrations between 50 and 500 μ M, substantially lower than the concentrations used for *B. xylophilus*. The survival of wild-type *C. elegans* N2 decreased significantly ($p < 0.05$) with increasing concentrations of H_2O_2 (Fig 6). Interestingly, both of the transgenic *C. elegans* overexpressing *B. xylophilus* catalases were resistant to H_2O_2 stress conditions. KHA149 and KHA151, expressing *Bxy-ctl-1* and *Bxy-ctl-2* respectively, were not statistically different ($p > 0.05$) in the H_2O_2 conditions tested.

Discussion

Depending on their lifestyle, sedentary or migratory, plant parasitic nematodes possess different repertoires of enzymatic and non-enzymatic molecules involved in oxidative stress metabolism that are a reflection of their pathogenicity [15]. In some organisms, like bacteria, these enzymes are defined as virulence factors [32]. Recent work [18] indicates that *B. xylophilus* has an enzymatic inventory to counterattack the pine oxidative burst. A total of 12 anti-oxidant enzymes were found in the *B. xylophilus* secretome, including PRX, CTL, GST, SOD, nucleoredoxin-like protein and thioredoxin [19]. In the present study, we investigated the tolerance of *B. xylophilus* isolates with different levels of virulence (high and low virulence) under *in vitro* oxidative stress conditions using H_2O_2 as an oxidizing agent. We were able to observe a contrasting performance between high and low virulence *B. xylophilus* and its relative *B. mucronatus*, corroborating previous results in other plant parasitic nematodes [33–34]. Emphasizing

A. KHA149 {*unc-119(ed3)III*; *chuEx149[Pctl-1::Bxy-ctl-1::gfp, pDP#MM016B]*}



B. KHA151 {*unc-119(ed3)III*; *chuEx151[Pctl-3::Bxy-ctl-2::gfp, pDP#MM016B]*}




Fig 5. Expression patterns of *Bxy-ctl-1::gfp* (A) and *Bxy-ctl-2::gfp* (B) in, respectively, transgenic *C. elegans* KHA149 and KHA151. Differential interference contrast (DIC) microscope images and, DIC and fluorescence-merged images (DIC+FL) of *C. elegans* head, vulva and tail region. Scale bars, 100 μ m.

doi:10.1371/journal.pone.0123839.g005

the results of the oxidative stress tolerance and H_2O_2 neutralization tests, the CTL (*Bxy-ctl-1* and *Bxy-ctl-2*) gene expression indicated that high and low virulence *B. xylophilus* respond differently towards H_2O_2 stress. High virulence nematodes showed an up-regulation of CTL genes under induced oxidative stress conditions, corroborating our preliminary results [24]. In addition, since all *B. xylophilus* isolates share protein sequence similarity between CTLs, the contrasting responses to H_2O_2 stress may indicate different regulation in the detoxification of ROS. This correlation between catalase levels and *in vitro* H_2O_2 tolerance has also been described in other parasitic nematodes [35–36]. Previously, Shinya et al. [20] showed the accumulation of anti-oxidant and detoxifying enzymes on the body surface of the nematode. Here, we were able to determine that both *B. xylophilus* catalases (*Bxy-ctl-1* and *Bxy-ctl-2*) are expressed in the intestine.




Fig 6. Survival percentage of wild-type (N2) and transgenic (KHA149 and KHA151) *C. elegans* after 24 hours exposition to H₂O₂ (H₂O₂ concentrations ranging between 0 and 500 μM). Error bars represent standard deviation. Different letters above the columns indicate significant differences ($p < 0.05$) between *C. elegans* N2, KHA149 and KHA151 survival percentages in each H₂O₂ treatment, according to post-hoc Duncan's test.

doi:10.1371/journal.pone.0123839.g006

We also accounted for other antioxidant enzymes (SODs and GXP) under H₂O₂ stress (S1 Fig), and found that, in our experimental conditions, only catalase expressions were significant. Although we could ascertain the importance of CTLs (*Bxy-ctl-1* and *Bxy-ctl-2*) in the protection against H₂O₂ induced-stress, we cannot rule out the effect of other antioxidant proteins in their response. Another important observation in this study is the maintenance of CTL induction in high virulence *B. xylophilus* even after 24h of H₂O₂ exposure. Kotze [37] observed similar CTL induction in parasitic nematode *Haemonchus contortus* against *in vitro* H₂O₂ stress. This feature is crucial and highlights the adaptive response and antioxidant protection of high virulence *B. xylophilus* in host conditions. Further studies are still needed to completely understand the role of all protectant molecules (enzymatic or non-enzymatic) of *B. xylophilus* against the massive ROS production of susceptible pines.

The regulatory pathways of xenobiotic degradation of *C. elegans* are conserved in *B. xylophilus* [18]. Hence, We constructed transgenic *C. elegans* overexpressing both *Bxy-ctl-1* and *Bxy-ctl-2*. Since we previously have shown that *Bxy-ctl-1* sequence was more similar to *Ce-ctl-1* and *Ce-ctl-2*, and *Bxy-ctl-2* to *Ce-ctl-3* [24], we chose *C. elegans Pctl-1* to regulate *Bxy-ctl-1* and *C. elegans Pctl-3* to regulate *Bxt-ctl-2*. Our results showed that subcellular localization of both catalases is in accordance with *C. elegans ctl-1* and *ctl-3* promoter expression as shown by Petriv and Rachubinski [38]. Furthermore, Oláhova et al. [31] have also shown a decrease in survival rate of wild-type N2 to 1 mM H₂O₂ only 10 hours after treatment, which support our selection to test 50–500 μM H₂O₂ as experimental concentrations. *C. elegans* contains three catalases, *Ce-ctl-1* (cytosolic), *Ce-ctl-2* (peroxisomal) and *Ce-ctl-3* (neurons, muscles and pharynx), which present different activities, with CTL-2 being responsible for 80% of total catalase activity [31]. Expressing exogenous *B. xylophilus* catalases, transgenic KHA149 and KHA151 animals were able to withstand the H₂O₂ stress up to 500 μM, supporting the efficiency of *B. xylophilus* catalases in H₂O₂ neutralization.

Supporting Information

S1 Fig. Relative gene expression of SODs (*Bxy-sod-1*; *Bxy-sod-2*; *Bxy-sod-3*) and GXP (*Bxy-gxp-1*; *Bxy-gxp-2*; *Bxy-gxp-3*) of *B. xylophilus* (high virulence isolates Ka4 and T4; low virulence isolate C14-5) under oxidative stress conditions. Error bars represent standard deviation. (TIF)

Acknowledgments

B. xylophilus isolates were provided by FFPRI, Tsukuba, Japan. *C. elegans* strains were provided by the *Caenorhabditis* Genetics Center (University of Minnesota, Minneapolis, MN). The authors thank to Margarida Espada (NemaLab, Universidade de Évora, Évora, Portugal) for the support in the *In-situ* hybridization experiments; and Dr. Mihoko Kato for linguistic correction of the manuscript.

Author Contributions

Conceived and designed the experiments: CSLV KH. Performed the experiments: CSLV YI KH. Analyzed the data: CSLV RS KH. Contributed reagents/materials/analysis tools: KH MM. Wrote the paper: CSLV KH. Read and approved the final manuscript: CSLV YI RS MM KH.

References

1. Futai K. Pine wood nematode, *Bursaphelenchus xylophilus*. *Ann Rev Phytopathol*. 2013; 51: 61–83. doi: [10.1146/annurev-phyto-081211-172910](https://doi.org/10.1146/annurev-phyto-081211-172910) PMID: [23663004](https://pubmed.ncbi.nlm.nih.gov/23663004/)
2. Mota MM, Vieira P. Pine Wilt Disease: A Worldwide Threat to Forest Ecosystems. 2008. Netherlands: Springer.
3. Vicente C, Espada M, Vieira P, Mota M. Pine Wilt Disease: a threat to European forestry. *Eur J Plant Pathol*. 2012; 133: 89–99.
4. Shinya R, Takeuchi Y, Ichimura K, Takemoto S, Futai K. Establishment of a set of inbred strains of the pine wood nematode, *Bursaphelenchus xylophilus* (Aphelenchida: Aphelenchoididae), and evidence of their varying levels of virulence. *Appl Entomol Zool*. 2012; 47: 341–350.
5. Kiyohara T, Bolla RI. Pathogenic variability among populations of the pinewood nematode, *Bursaphelenchus xylophilus*. *Forest Sci*. 1990; 36: 1061–1076.
6. Aikawa T, Kikuchi T. Estimation of virulence of *Bursaphelenchus xylophilus* (Nematoda: Aphelenchoididae) based on its reproductive ability. *Nematol*. 2007; 9: 371–377.
7. Asai E, Futai K. Effects of inoculum density of pinewood nematode on the development of pine wilt disease in Japanese black pine seedlings pretreated with simulated acid rain. *Forest Pathol*. 2005; 35: 135–144.
8. Aikawa T, Togashi K, Kosaka H. Different developmental responses of virulent and avirulent isolates of the pine wood nematode, *Bursaphelenchus xylophilus* (Nematoda: Aphelenchoididae), to the insect vector, *Monochamus alternatus* (Coleoptera: Cerambycidae). *Environm Entomol*. 2003; 32: 96–102.
9. Takemoto S. Population ecology of *Bursaphelenchus xylophilus*. In: *Pine Wilt Disease*. 2008. Netherlands: Springer.
10. D'Autréaux B, Toledano MB. ROS as signalling molecules: mechanisms that generate specificity in ROS homeostasis. *Nat Rev Mol Cell Biol*. 2007; 10: 813–824. PMID: [17848967](https://pubmed.ncbi.nlm.nih.gov/17848967/)
11. Quan LJ, Zhang B, Shi WS, Li HY. Hydrogen peroxide in plants: a versatile molecule of the reactive oxygen species network. *J Integrative Plant Biol*. 2008; 50: 2–18.
12. Lamb C, Dixon R. The oxidative burst in plant disease resistance. *Annu Rev Plant Physiol Plant Mol Biol*. 1997; 48: 251–275. PMID: [15012264](https://pubmed.ncbi.nlm.nih.gov/15012264/)
13. Dubreuil G, Deleury E, Magliano M, Jaouannet M, Abad P, et al. Peroxiredoxins from the plant parasitic root-knot nematode, *Meloidogyne incognita*, are required for successful development within the host. *Int J Parasitol*. 2011; 41: 385–396. doi: [10.1016/j.ijpara.2010.10.008](https://doi.org/10.1016/j.ijpara.2010.10.008) PMID: [21145323](https://pubmed.ncbi.nlm.nih.gov/21145323/)
14. Henkle-Dührsen K, Kampkötter A. Antioxidant enzyme families in parasitic nematodes. *Mol Biochem Parasitol*. 2001; 114: 129–142. PMID: [11378193](https://pubmed.ncbi.nlm.nih.gov/11378193/)

15. Molinari S, Miacola. Antioxidant enzymes in phytoparasitic nematodes. *J Nematol.* 1997; 29: 153–159. PMID: [19274144](#)
16. Robertson L, Robertson WM, Sobczak M, Helder J, Tetaud E, et al. Cloning, expression and functional characterisation of a peroxiredoxin from the potato cyst nematode *Globodera rostochiensis*. *Mol Biochem Parasitol.* 2000; 111: 41–49. PMID: [11087915](#)
17. Bellafiore S, Shen Z, Rosso MN, Abad P, Shih P, et al. Direct identification of the *Meloidogyne incognita* secretome reveals proteins with host cell reprogramming potential. *PLoS pathogens.* 2008; 4: e1000192. doi: [10.1371/journal.ppat.1000192](#) PMID: [18974830](#)
18. Kikuchi T, Cotton JA, Dalzell JJ, Hasegawa K, Kanzaki N, et al. Genomic insights into the origin of parasitism in the emerging plant pathogen *Bursaphelenchus xylophilus*. *PLoS Pathogens.* 2011; 7:e 1002219.
19. Shinya R, Morisaka H, Kikuchi T, Takeuchi Y, Ueda M, et al. Secretome analysis of pine wood nematode *Bursaphelenchus xylophilus* reveals the tangled roots of parasitism and its potential for molecular mimicry. *PloS one.* 2012; 8:e 67377.
20. Shinya R, Morisaka H, Takeuchi Y, Ueda M, Futai K. Comparison of the surface coat proteins of the pine wood nematode appeared during host pine infection and in vitro culture by a proteomic approach. *Phytopathol.* 2010; 100: 1289–97.
21. Hirao T, Fukatsu E, Watanabe A. Characterization of resistance to pine wood nematode infection in *Pinus thunbergii* using suppression subtractive hybridization. *BMC Plant Biol.* 2012; 12:13. doi: [10.1186/1471-2229-12-13](#) PMID: [22272988](#)
22. Santos CSS, Vasconcelos MW. Identification of genes differentially expressed in *Pinus pinaster* and *Pinus pinea* after infection with pine wood nematode. *Eur J Plant Pathol.* 2012; 132: 407–418.
23. He L, Wu X, Yu L, Ji J, Ye J. The difference of H₂O₂ and oxidative enzyme in the interaction of different resistance pines and *Bursaphelenchus xylophilus*. *J Nanjing Univ Natur Sci.* 2010; 06.
24. Vicente CSL, Ikuyo Y, Mota M, Hasegawa K. Pinewood nematode-associated bacteria contribute to oxidative stress resistance of *Bursaphelenchus xylophilus*. *BMC Microbiol.* 2013; 13: 299. doi: [10.1186/1471-2180-13-299](#) PMID: [24365493](#)
25. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔCT} method. *Methods.* 2001; 25: 402–408. PMID: [11846609](#)
26. Rozen S, Skaletzki H. Primer3 on the www for general users and for biologist programmers. *In: Bioinformatics Methods and Protocols: Methods in Molecular Biology* Totowa. 2000; Humana Press.
27. Petersen TN, Brunak S, Heijne G, Nielsen H. SignalP 4.0: discriminating signal peptides from transmembrane regions *Nat Methods.* 2011; 8:785–786. doi: [10.1038/nmeth.1701](#) PMID: [21959131](#)
28. De Boer JM, Yan Y, Smant G, Davis EL, Baum TJ. In-situ hybridization to messenger RNA in *Heterodera glycines*. *J. Nematol.* 1998; 30: 309–312. PMID: [19274224](#)
29. Kikuchi T, Jones JT, Aikawa T, Kosaka H, Ogura N. A family of glycosyl hydrolase family 45 cellulases from pine wood nematode *Bursaphelenchus xylophilus*. *FEBS letters.* 2004; 572: 201–205. PMID: [15304348](#)
30. Brenner S. The genetics of the nematode *Caenorhabditis elegans*. *Genetics.* 1974; 77: 71–94. PMID: [4366476](#)
31. Oláhova M, Taylor SR, Khazaipoul S, Wang J, Morgan BA, et al. A redox-sensitive peroxiredoxin that is important for longevity has tissue- and stress-specific roles in stress resistance. *PNAS.* 2008; 50: 19839–19844. doi: [10.1073/pnas.0805507105](#) PMID: [19064914](#)
32. Guo M, Block A, Bryan CD, Becker DF, Alfano AJ. *Pseudomonas syringae* catalases are collectively required for plant pathogenesis. *J Bacteriol.* 2012; 194: 5054–5064. doi: [10.1128/JB.00999-12](#) PMID: [22797762](#)
33. Molinari S, Rosso L, Ornat LC. The role of antioxidant enzymes in the virulence of root-knot nematodes on resistant tomato. *Nematopica.* 2005; 35: 88.
34. Molinari S. Antioxidant enzymes in (a) virulent populations of root-knot nematodes. *Nematol.* 2009; 11: 689–697.
35. Ou X, Thomas GR, Chacon MR, Tang L, Selkirk ME. *Brugia malayi*: differential susceptibility to and metabolism of hydrogen peroxide in adults and microfilariae. *Exp Parasitol.* 1995; 80: 530–540. PMID: [7729488](#)
36. Kotze AC, McClure SJ. *Haemonchus contortus* utilises catalase in defence against exogenous hydrogen peroxide in vitro. *Int J Parasitol.* 2001; 31: 1563–1571. PMID: [11730782](#)
37. Kotze AC. Catalase induction protects *Haemonchus contortus* against hydrogen peroxide in vitro. *Int J Parasitol.* 2003; 33: 393–400. PMID: [12705932](#)
38. Petriv OI, Rachubinski RA. Lack of peroxisomal catalase causes a progeric phenotype in *Caenorhabditis elegans*. *J Biol Chem.* 2004; 279: 19996–20001. PMID: [14996832](#)